

- for liver fibrosis assessment [J]. *Ultrasound Med Biol*, 2013, 39 (11): 1933 - 1941.
- [41] ZENG J, ZHENG J, HUANG ZP, et al. The comparative study of two-dimensional shear wave elastography for assessing hepatic fibrosis in chronic hepatitis B using different taking value methods [J]. *Chin J Ultras Med*, 2016, 32(8): 717 - 720. (in Chinese)
曾婕, 郑剑, 黄泽萍, 等. 二维剪切波弹性成像诊断慢性乙型肝炎肝纤维化不同取值方法的对比研究 [J]. *中国超声医学杂志*, 2016, 32(8): 717 - 720.
- [42] POYNARD T, MUNTEANU M, LUCKINA E, et al. Liver fibrosis evaluation using real-time shear wave elastography: applicability and diagnostic performance using methods without a gold standard [J]. *J Hepatol*, 2013, 58(5): 928 - 935.
- [43] YOON JH, LEE JM, JOO I, et al. Hepatic fibrosis: prospective comparison of MR elastography and US shear-wave elastography for evaluation [J]. *Radiology*, 2014, 273(3): 772 - 782.
- [44] ZHUANG Y, DING H, ZHANG Y, et al. Two-dimensional shear-wave elastography performance in the noninvasive evaluation of liver fibrosis in patients with chronic hepatitis B: comparison with serum fibrosis indexes [J]. *Radiology*, 2017, 283(3): 873 - 882.
- [45] PISCAGLIA F, SALVATORE V, MULAZZANI L, et al. Ultrasound shear wave elastography for liver disease. A critical appraisal of the many actors on the stage [J]. *Ultraschall Med*, 2016, 37 (1): 1 - 5.
- [46] COCO B, OLIVERI F, MAINA AM, et al. Transient elastography: a new surrogate marker of liver fibrosis influenced by major changes of transaminases [J]. *J Viral Hepat*, 2007, 14(5): 360 - 369.
- [47] MILLONIG G, REIMANN FM, FRIEDRICH S, et al. Extrahepatic cholestasis increases liver stiffness (FibroScan) irrespective of fibrosis [J]. *Hepatology*, 2008, 48(5): 1718 - 1723.
- [48] MILLONIG G, FRIEDRICH S, ADOLF S, et al. Liver stiffness is directly influenced by central venous pressure [J]. *J Hepatol Elsevier*, 2010, 52(2): 206 - 210.
- [49] BARDOU - JACQUET E, LEGROS L, SORO D, et al. Effect of alcohol consumption on liver stiffness measured by transient elastography [J]. *World J Gastroenterol*, 2013, 19(4): 516 - 522.

[本文首次发表于中华超声影像学杂志, 2017, 26 (11): 921 - 927]

引证本文: Panel of Elastography Assessment of Liver Fibrosis, Study Group of Interventional Ultrasound, Society of Ultrasound in Medicine of Chinese Medical Association. Guidelines for clinical application of two-dimensional shear wave elastography in assessment of liver fibrosis in chronic hepatitis B [J]. *J Clin Hepatol*, 2018, 34 (2): 255 - 261. (in Chinese)
中华医学会超声医学分会介入超声学组弹性成像评估肝纤维化专家组. 二维剪切波弹性成像评估慢性乙型肝炎肝纤维化临床应用指南 [J]. *临床肝胆病杂志*, 2018, 34(2): 255 - 261.

(本文编辑: 王亚南)

• 消息 •

《临床肝胆病杂志》免费刊发肝胆胰诊疗指南及指南译文

标准治疗指南是有助于医生或患者在特定临床条件下确定恰当医疗保健决定的系统性陈述(规定)。是把临床的思维方式,以一个系统化的文件固化下来,让非专业或者专业人士,直接参考它来处理面临的患者群体。

由于指南具有很大实用价值,本刊决定对国内外最新制订的肝胆胰指南及指南译文免收版面费并优先发表。

欢迎各组织机构和作者推荐指南(含指南,共识,规范等),推荐人请先与编辑部联系,经审核符合本刊刊载范围的方予以刊登。

本刊希望此举对于促进肝胆胰疾病的规范治疗,减少医疗纠纷和因为不正确治疗而造成的医源性伤害,产生积极的推动作用。

《临床肝胆病杂志》编辑部

2018年2月20日